

Imaging the quantal substructure of single IP₃R channel activity during Ca²⁺ puffs in intact mammalian cells

Ian F. Smith^{a,1} and Ian Parker^{a,b}

Departments of ^aNeurobiology and Behavior and ^bPhysiology and Biophysics, University of California at Irvine, Irvine, CA 92697-4550

Edited by Richard W. Aldrich, The University of Texas, Austin, TX, and approved February 20, 2009 (received for review October 27, 2008)

The spatiotemporal patterning of Ca²⁺ signals regulates numerous cellular functions, and is determined by the functional properties and spatial clustering of inositol trisphosphate receptor (IP₃R) Ca²⁺ release channels in the endoplasmic reticulum membrane. However, studies at the single-channel level have been hampered because IP₃Rs are inaccessible to patch-clamp recording in intact cells, and because excised organelle and bilayer reconstitution systems disrupt the Ca²⁺-induced Ca²⁺ release (CICR) process that mediates channel-channel coordination. We introduce here the use of total internal reflection fluorescence microscopy to image single-channel Ca²⁺ flux through individual and clustered IP₃R in intact mammalian cells. This enables a quantal dissection of the local calcium puffs that constitute building blocks of cellular Ca²⁺ signals, revealing stochastic recruitment of, on average, approximately 6 active IP₃R clusters within <500 nm. Channel openings are rapidly (≈10 ms) recruited by opening of an initial trigger channel, and a similarly rapid inhibitory process terminates puffs despite local [Ca²⁺] elevation that would otherwise sustain Ca²⁺-induced Ca²⁺ release indefinitely. Minimally invasive, nano-scale Ca²⁺ imaging provides a powerful tool for the functional study of intracellular Ca²⁺ release channels while maintaining the native architecture and dynamic interactions essential for discrete and selective cell signaling.

calcium signaling | single-channel flux | TIRF microscopy | optical patch-clamp

Cytosolic signals resulting from Ca²⁺ liberation from the endoplasmic reticulum (ER) through inositol trisphosphate receptor/channels (IP₃R) regulate cellular functions as diverse as gene expression, secretion, and synaptic plasticity (1). Information is encoded by the spatiotemporal patterning of cytosolic Ca²⁺ transients, which are organized in a hierarchical manner resulting from the clustering of IP₃R in the ER membrane. Local, “elementary” Ca²⁺ transients (i.e., Ca²⁺ puffs) are generated by the concerted openings of multiple IP₃R within a cluster, and serve autonomous signaling functions as well as constituting the building blocks from which global cellular Ca²⁺ waves are constructed (2–6). In turn, puffs are composed from “fundamental” signals (i.e., Ca²⁺ blips), representing Ca²⁺ flux through individual IP₃R (7–10). Information concerning the spatial localization and numbers of IP₃R involved in a puff, and the mechanisms and kinetics by which their activity is coordinated, is essential to understand how these channels act to initiate and terminate local Ca²⁺ liberation (11). However, studies at the single-channel level have been hampered because the intracellular location of IP₃R renders them inaccessible to patch-clamp recording within intact cells. Electrophysiological recordings have thus been restricted to excised nuclei or lipid bilayer reconstitution systems (11), approaches that provide little or no spatial information and disrupt the process of Ca²⁺-induced Ca²⁺ release (CICR) that mediates channel-channel interactions (12).

Our knowledge of local cellular Ca²⁺ signaling instead derives largely from minimally invasive whole-cell Ca²⁺ imaging tech-

niques. Nonetheless, recordings by conventional wide-field or confocal fluorescence microscopy fail to provide resolution at the single-channel level, and puffs imaged by these techniques show an apparently monotonic rising phase lasting tens of milliseconds as Ca²⁺ is liberated through open channels, followed by a slower decay as the Ca²⁺ micro-domain at the puff site dissipates following channel closure (4–6). Although specific instances of single IP₃R channel fluorescence signals have been reported as isolated blips (7, 13) and as “trigger” events immediately preceding puffs (10), it had not been possible to directly dissect the contributions of individual channels during puffs. Here, we capitalize on experimental and theoretical findings that the resolution of local Ca²⁺ signals can be dramatically improved by monitoring fluorescence from attoliter volumes around open channels by means of total internal reflection fluorescence (TIRF) microscopy (14). We had previously used this approach to image single-channel calcium fluorescence transients arising from gating of Ca²⁺-permeable channels in the plasma membrane (15, 16). Following the discovery that a population of puff sites in SH-SY5Y neuroblastoma cells lie sufficiently close to the plasma membrane to be within range of the approximately 100-nm evanescent field of the TIRF microscope (17), we now apply this technique to resolve the properties of individual intracellular IP₃R channels, and to dissect their contributions during Ca²⁺ puffs within intact mammalian cells.

Results

Resolving Ca²⁺ Fluorescence Signals from Individual IP₃Rs. We loaded SH-SY5Y cells with fluo-4 and caged iIP₃ by incubation with membrane-permeant esters, and imaged local Ca²⁺ puffs evoked by photo-released i-IP₃. Puffs imaged by wide-field epifluorescence showed smoothly rising and falling phases (Fig. 1A, gray trace), on which it was not possible to resolve the underlying step-wise changes that would be expected as individual channels stochastically open and close. To then achieve single-channel resolution, we used TIRF imaging of a subset of puff sites located close to the plasma membrane (17), so that rapid binding of Ca²⁺ to the fast indicator dye within attoliter cytosolic volumes around puff sites yields fluorescence signals that more closely track instantaneous Ca²⁺ flux through IP₃R channels (14). Additionally, we gained a further improvement by loading cells with EGTA, a Ca²⁺ buffer that inhibits wave propagation (17–19) and accelerates the collapse of the local Ca²⁺ micro-domain but, because of its slow binding kinetics, minimally perturbs local free [Ca²⁺] within a cluster (14) and has little effect on peak puff amplitudes (17–19). EGTA is thus

Author contributions: I.F.S. and I.P. designed research, performed research, analyzed data, and wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

¹To whom correspondence should be addressed. E-mail: ismith@uci.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0810799106/DCSupplemental.




Fig. 3. Ca^{2+} sources during different amplitude steps at a puff site localize within a few hundred nanometers. (A) Ca^{2+} signal from a $1\text{-}\mu\text{m}^2$ region of interest showing 5 discrete amplitude levels during a single puff. (B) Sequence of images showing the location of the Ca^{2+} fluorescence at times corresponding to the numbered levels in A. Each panel is an average of image frames captured during that step, and cross-hairs mark the centroid position of the fluorescence signal in the first frame. (C) Extended record showing multiple events at the same region of interest. The puff marked by the asterisk is that illustrated in A. (D) Scatter plot marks the centroid positions of Ca^{2+} signals during all ($n = 53$) step amplitude levels throughout the record in C, derived by fitting 2D Gaussian functions to averaged images like those in B. Square marks the region of interest from which the fluorescence traces in A and B were measured. The curve shows, on the same scale, the spatial distribution of fluorescence measured along a line passing through the center of the puff in B1.

amplitude of blips with that ($\Delta F/F_0 \approx 0.5$) of single nicotinic receptors passing a Ca^{2+} current of approximately 0.25 pA (16) implies a Ca^{2+} current of approximately 0.05 pA , comparable to that ($\approx 0.1\text{ pA}$) estimated for IP_3R channels in the intact cell (11).

Quantal Analysis of Ca^{2+} Puffs. Importantly, puffs imaged by TIRF microscopy showed abrupt step-wise transitions between fluorescence levels (Fig. 2C and Movie S2), predominantly during their falling phase. We can exclude that these steps arose because the site of Ca^{2+} liberation shifted outside the region of interest to cause artifactually smaller fluorescence signals as Ca^{2+} became diluted by diffusion. Instead, the centroid locations of Ca^{2+} -fluorescence signals during multiple steps deviated by no more than few hundred nanometers of one another (Fig. 3 and Movie S3), a distance within the region of interest used to measure the traces, and small in comparison to the spatial spread of the fluorescence signal (Fig. 3D). We thus conclude that transitions in amplitude levels arise from the step-wise recruitment and closing of varying numbers of IP_3Rs , localized within clusters with dimensions of $<500\text{ nm}$.

This concept is further supported by a quantal analysis (23) of amplitude levels, showing a multimodal distribution with recurring peaks at integer multiples of the unitary blip amplitude (Fig. 4A). Although it has been proposed that local depletion of ER Ca^{2+} may result in an incrementally smaller single-channel Ca^{2+} flux as progressively greater numbers of IP_3R channels open (24), our results suggest that the fluorescence signals contributed by each channel summate linearly. Discrete peaks remain evident in the step amplitude distribution at equal increments as great as 6 or 7 times the unitary amplitude (Fig. 4A); analysis of step decrements from the peak of large puffs (>15 times the unitary event) did not reveal steps appreciably smaller than the unitary event (data not shown), and even the largest signals ($\Delta F/F_0 \approx 2$) are well within the linear range of Fluo-4 ($F_{\text{max}}/F_{\text{min}} > 20$). We thus estimate the number of IP_3R channels open at a site at any given time simply by dividing the fluorescence signal ($\Delta F/F_0$) by the mean unitary event signal at that site.

Intra- and Inter-Site Variability in Numbers of IP_3Rs . The numbers of channels open at the peak of each event follow a highly skewed distribution, with many events involving concurrent openings of only 1 or 2 channels and progressively fewer involving as many as 20 channels (Fig. 4B). Much of this variability results from inter-site heterogeneity, as shown in Fig. 4C, plotting the amplitude distribution of the largest event seen at any given site. Because the photolysis flash typically evoked tens of events at each site, this provides an approximation of the minimal total number of functional IP_3R channels within that cluster. The mean number per site is 6.02 ± 0.48 ($n > 80$ sites), and again follows a highly skewed distribution, including a few ($\approx 6\%$) sites comprising a lone channel, a majority of sites comprising a handful of channels, and a small proportion ($\approx 16\%$) containing >10 channels. These data pool observations from many cells, but observations of widely varying maximal event amplitudes among sites within an individual cell (e.g., $\Delta F/F_0$, 0.15–2.26) indicate that the variability arises primarily from inter-site variability rather than from differences between cells. To then examine the variability among successive responses at a given site, we minimized inter-site variability by pooling data from selected sites with maximal event sizes 5 to 10 times the unitary signal. This revealed an approximately equal proportion of events involving as many as 6 channels, and decreasing numbers of larger events (Fig. 4D).

What might account for this wide variability in responses at a given puff site? We considered several simplified models, based on the assumption that CICR evoked by spontaneous opening of an initial channel triggers a puff by increasing the opening probability of neighboring IP_3Rs in the cluster (25). Fig. 4E plots the data in terms of the number of additional channels responding to this trigger. We first considered a model comprised of n independent channels, each of which has an equal probability p of being triggered. The open symbols in Fig. 4E shows the predicted distribution for $n = 10$, and with $P = 0.34$ so as to match the observed mean number of responding channels. This provides a poor fit to the data, and other simulations with between 8 and 50 channels similarly failed to account for the




Fig. 5. IP₃R channel gating kinetics during blips and puffs. (A) Distribution of unitary blip event durations, after excluding the small proportion of events with durations >100 ms. The curve is a single exponential with time constant τ of 17 ms. (B) Histogram shows the distribution of puff durations, measured as the interval between attainment of peak amplitude and return to baseline and expressed relative to the mean blip duration of 17 ms. Data were selected from events with normalized peak amplitudes between 5 and 10, and exclude events that showed channel openings during the falling phase. The black curve shows simulated data generated assuming that 8 channels are open at the peak of the puff, and then close stochastically and independently after a mean dwell time of 17 ms. (C) Mean time to event termination as a function of the estimated number of channels open at the peak. Open symbols are experimental data. Filled symbols show the relationship predicted if channels close stochastically and independently after a mean lifetime of 17 ms. (D and E) Graphs plot, respectively, the rates of channel openings/closings as functions of time after initiation of a puff. Symbols are individual measurements of numbers of channels that opened/closed during successive 5-ms time bins, estimated from the normalized fluorescence change during that interval (Left). Curves show the mean rates of channel opening/closing averaged over varying time bins (Right).

initial positive feedback, it is surprising that puffs are not stereotypical, all-or-none events, but rather show great variability such that, on average, only approximately half of the available IP₃R channels open (Fig. 4D). This is unlikely to result in stochastic recovery of IP₃R from an inactivated state induced by a preceding puff, because the mean size of the first puffs evoked by photo-released IP₃ is not appreciably greater than subsequent puffs (17), and puff amplitudes show only weak correlations with the amplitude of a preceding puff and with the inter-puff interval (31). Instead, we propose that the rapid cessation of channel openings results from an inhibitory process with kinetics comparable to the regenerative activation. The variability in puff amplitudes (peak number of open channels) at a given site may then reflect a stochastic interplay between activation and inactivation processes, such that an appreciable proportion of IP₃R become inactivated before they have a chance to open.

The mechanism of termination of puffs (and analogous sparks mediated by RyRs) is of great interest, because CICR is otherwise expected to prolong these events indefinitely (32, 33). Among several postulated mechanisms (33), our results rule out stochastic attrition (i.e., breakdown of CICR because all channels randomly happen to be closed at the same time) as the sole explanation, as this predicts that puff durations will increase exponentially with the number of open channels (30). Moreover, the step-wise decay of puffs at integer quantal levels indicates that termination results from channel gating and not simply because local exhaustion of ER Ca²⁺ stores diminishes the single-channel Ca²⁺ flux. Instead, the progressive closure of IP₃R channels after the peak of a puff, with relatively few subsequent openings, is consistent with the rapid onset of a

strong inhibitory process as discussed earlier. Our data do not directly address the nature of this inhibition, but a likely candidate is the strong Ca²⁺-dependent inhibition exhibited by the type 1 IP₃R predominant in this cell type. Channels that are open at the peak of the puff subsequently close after a mean duration characteristic of the single-channel lifetime, and infrequent openings on the falling phase may reflect slow recovery of inactivated channels.

In conclusion, IP₃-mediated Ca²⁺ signals are an example in which stochastic fluctuations at the single-channel level give rise to whole-cell responses (11). Development of minimally invasive, functional single-channel imaging methods thus represents a major advance in elucidating the link between these nanoscopic and macroscopic cell signaling processes. By its nature, TIRF microscopy is likely to be limited to cultured or isolated cells in which the ER/sarcoplasmic reticulum is close to the cell membrane, and may thus be unable to fully reveal potential complexities of channel clustering and IP₃ gradients within native tissues. Nevertheless, our approach should be widely applicable to studies of the functional architecture and dynamics of intracellular ion channels in intact cells at the truly single-molecule level.

Materials and Methods

Cell Culture and Loading. Human neuroblastoma SH-SY5Y cells were cultured as previously described (17) in a mixture (1:1) of Ham's F12 medium and Eagle MEM, supplemented with 10% (vol./vol.) FCS and 1% nonessential amino acids. Cells were incubated at 37 °C in a humidified incubator gassed with 95% air and 5% CO₂, passaged every 7 days, and used for a maximum of 20 passages. Four days before imaging, cells were harvested in PBS solution without Ca²⁺ or Mg²⁺ and subcultured in Petri dishes with glass coverslips as

